

### *Uromycladium tepperianum* on *Acacia* spp.

*Uromycladium tepperianum* is a microcyclic rust that infects more than one hundred species of *Acacia* (Gathe 1971) and several other genera in the Fabaceae, causing large, conspicuous galls (Morris 1987). *Acacia pycnantha* Benth., cultivated in Australia for its bark, is severely affected by *U. tepperianum*, which causes significant yield losses and eventually the death of the host (Gathe 1971). However, this rust has potential as a biocontrol agent for weedy acacias outside of Australia, for example, *U. tepperianum* has been proven a highly effective against *A. saligna* in South Africa (Morris 1997, Wood & Morris 2007).

*Uromycladium tepperianum* (Sacc.) McAlpine, Ann. Mycol. 3: 310. 1905.

**Spermogonia** minute, brownish then black, globose, 150 µm diam; spermatia hyaline, ellipsoid.

**Aecia** and **uredinia** unknown.

**Telia** develop on galls on leaves, branches, inflorescences, and fruits; infections causing swollen, distorted galls up to 18 × 6 cm, and witches' brooms of different shapes and sizes, cinnamon to chocolate brown, powdery; **teliospores** composed of a cluster of three probasidial cells at top of a single pedicel, depressed globose to globose, cinnamon brown, thickly vertically striate, margin crenulate, wall 2-3 µm, at apex up to 5 µm thick, 14-22 µm high, 18-25 µm wide, one apical germ pore; pedicel hyaline, septate, deciduous.

**Hosts:** Species of *Acacia*, *Albizia* and *Racosperma* (Fabaceae)

**Geographic Distribution** Australia, Java, New Caledonia, New Zealand, Papua New Guinea, and South Africa.

*Uromycladium* is characterized by the production of teliospores composed of 1-3 probasidial cells, with or without cysts, on a single pedicel. Eight species of *Uromycladium* have been described occurring primarily on *Acacia* in Australia and New Zealand. Teliospores of *Uromycladium tepperianum* and *U. notabile* McAlpine have three probasidial cells and no cysts (Burgess 1934). *Uromycladium notabile* produces uredinia, and the probasidial cells have linear verrucae, while no uredinia are known for *U. tepperianum* and the probasidial cells are distinctly striate. *Uromycladium acaciae* (Cooke) P. Syd. & Syd. produces teliospores with two probasidial cells and no cysts. Teliospores of *Uromycladium simplex* McAlpine and *Uromycladium robinsoni* McAlpine have one globose probasidial cell and one cyst. *Uromycladium fusisporum* (Cooke & Massee) Savile has teliospores with one probasidial cells and no cyst (Savile 1971). *Uromycladium maritimum* McAlpine and *Uromycladium alpinum* McAlpine produce teliospores with two probasidial cells and one cyst.

#### References:

- Burgess, A.** 1934. Studies in the genus *Uromycladium* (Uredinaceae) I. Proc. Linn. Soc. New South Wales 59: 212-228.
- Gathe, J.** 1971. Host range and symptoms in western Australia of gall rust, *Uromycladium tepperianum*. J. Roy. Soc. W. Australia 54: 114-118.
- McAlpine, D.** 1906. The Rusts of Australia. Their Structure, Nature and Classification. Department of Agriculture, Victoria, 349 pages.
- Morris, M.J.** 1987. Biology of the *Acacia* gall rust, *Uromycladium tepperianum*. Pl. Pathol. 36: 100-106.
- Morris, M.J.** 1991. The use of plant pathogens for biological weed control in South Africa. Agric. Eco-syst. Environm. 37: 239-255.
- Morris, M.J.** 1997. Impact of the gall-forming rust fungus *Uromycladium tepperianum* on the invasive tree *Acacia saligna* in South Africa. J. Biol. Control 10: 75-82.
- Morris, M.J.** 1999. The contribution of the gall-forming rust fungus *Uromycladium tepperianum* (Sacc.) McAlp. to the biological control of *Acacia saligna* (Labill.) Wendl. (Fabaceae) in South Africa. African Entomol. Mem. 1: 125-128.
- Savile, D.B.O.** 1971. Generic disposition and pycnium type in Uredinales. Mycologia 63: 1089-1091.
- Wood, A.R., and Morris, M.J.** 2007. Impact of the gall-forming rust fungus *Uromycladium tepperianum* on the invasive tree *Acacia saligna* in South Africa: 15 years of monitoring. Biol. Control 41: 68-77.

Telia BPI 871523



Telia BPI 871523



Telia BPI 871523




Telia BPI 871523



Telia BPI 871523



Teliospores BPI 871523




Teliospores BPI 871523




Teliospores BPI 871523



Teliospores BPI 871523



Teliospores BPI 871523



Teliospores BPI 871523

